

This Page Is Inserted by IFW Operations
and is not a part of the Official Record

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images may include (but are not limited to):

- BLACK BORDERS
- TEXT CUT OFF AT TOP, BOTTOM OR SIDES
- FADED TEXT
- ILLEGIBLE TEXT
- SKEWED/SLANTED IMAGES
- COLORED PHOTOS
- BLACK OR VERY BLACK AND WHITE DARK PHOTOS
- GRAY SCALE DOCUMENTS

IMAGES ARE BEST AVAILABLE COPY.

As rescanning documents *will not* correct images,
please do not report the images to the
Image Problem Mailbox.

THIS PAGE BLANK (USPTO)



The
Patent
Office

PCT/GB98 / 0 1 5 8 0

The Patent Office
Concept House
Cardiff Road
Newport
South Wales
NP9 1RH

PRIORITY DOCUMENT

SUBMITTED OR TRANSMITTED IN
COMPLIANCE WITH RULE 17.1(a) OR (b)

REC'D 0 3 JUL 1998

WIPO

PCT

I, the undersigned, being an officer duly authorised in accordance with Section 74(1) and (4) of the Deregulation & Contracting Out Act 1994, to sign and issue certificates on behalf of the Comptroller-General, hereby certify that annexed hereto is a true copy of the documents as originally filed in connection with the patent application identified therein.

In accordance with the Patents (Companies Re-registration) Rules 1982, if a company named in this certificate and any accompanying documents has re-registered under the Companies Act 1980 with the same name as that with which it was registered immediately before re-registration save for the substitution as, or inclusion as, the last part of the name of the words "public limited company" or their equivalents in Welsh, references to the name of the company in this certificate and any accompanying documents shall be treated as references to the name with which it is so re-registered.

In accordance with the rules, the words "public limited company" may be replaced by p.l.c., plc, P.L.C. or PLC.

Re-registration under the Companies Act does not constitute a new legal entity but merely subjects the company to certain additional company law rules.

Signed

Andrew Gersey

Dated 22nd June 1998

THIS PAGE BLANK (USPTO)

Request for grant of a patent

(See the notes on the back of this form. You can also get an explanatory leaflet from the Patent Office to help you fill in this form)



The Patent Office

Cardiff Road
Newport
Gwent NP9 1RH

1. Your reference

PA 419 A

2. Patent appli

(The Patent Of

9722674.0

27 OCT 1997

3. Full name, address and postcode of the or of each applicant (underline all surnames)

CELLTECH THERAPEUTICS LIMITED
216 BATH ROAD
SLOUGH, SL1 4EN

Patents ADP number (if you know it)

If the applicant is a corporate body, give the country/state of its incorporation

UK 6672877001

4. Title of the invention

CHEMICAL COMPOUNDS

5. Name of your agent (if you have one)

"Address for service" in the United Kingdom to which all correspondence should be sent (including the postcode)

PATENT DEPARTMENT
CELLTECH THERAPEUTICS LIMITED
216 BATH ROAD
SLOUGH
SL1 4EN

Patents ADP number (if you know it)

6672877002

6. If you are declaring priority from one or more earlier patent applications, give the country and the date of filing of the or of each of these earlier applications and (if you know it) the or each application number

Country

Priority application number
(if you know it)

Date of filing
(day / month / year)

7. If this application is divided or otherwise derived from an earlier UK application, give the number and the filing date of the earlier application

Number of earlier application

Date of filing
(day / month / year)

8. Is a statement of inventorship and of right to grant of a patent required in support of this request? (Answer 'Yes' if:

- a) any applicant named in part 3 is not an inventor, or
 - b) there is an inventor who is not named as an applicant, or
 - c) any named applicant is a corporate body.
- See note (d))

Yes

Patents Form 1/77

9. Enter the number of sheets for any of the following items you are filing with this form. Do not count copies of the same document

Continuation sheets of this form ☐

Description 32

Claim(s) ☐ 2

Abstract ☐

Drawing(s) ☐

10. If you are also filing any of the following, state how many against each item.

Priority documents ☐

Translations of priority documents ☐

Statement of inventorship and right to grant of a patent (Patents Form 7/77) ☐

Request for preliminary examination and search (Patents Form 9/77) ☐

Request for substantive examination (Patents Form 10/77) ☐

Any other documents ☐
(please specify)

11. I/We request the grant of a patent on the basis of this application.

Signature

[Signature]

Date

27th October 1

12. Name and daytime telephone number of person to contact in the United Kingdom

DR P. ANSELL 01753 534655

Warning

After an application for a patent has been filed, the Comptroller of the Patent Office will consider whether publication or communication of the invention should be prohibited or restricted under Section 22 of the Patents Act 1977. You will be informed if it is necessary to prohibit or restrict your invention in this way. Furthermore, if you live in the United Kingdom, Section 23 of the Patents Act 1977 stops you from applying for a patent abroad without first getting written permission from the Patent Office unless an application has been filed at least 6 weeks beforehand in the United Kingdom for a patent for the same invention and either no direction prohibiting publication or communication has been given, or any such direction has been revoked.

Notes

- If you need help to fill in this form or you have any questions, please contact the Patent Office on 0645 500505.
- Write your answers in capital letters using black ink or you may type them.
- If there is not enough space for all the relevant details on any part of this form, please continue on a separate sheet of paper and write "see continuation sheet" in the relevant part(s). Any continuation sheet should be attached to this form.
- If you have answered 'Yes' Patents Form 7/77 will need to be filed.
- Once you have filled in the form you must remember to sign and date it.
- For details of the fee and ways to pay please contact the Patent Office.

CHEMICAL COMPOUNDS

5 This invention relates to a series of tyrosine derivatives, to compositions containing them, to processes for their preparation, and to their use in medicine.

10 Over the last few years it has become increasingly clear that the physical interaction of inflammatory leukocytes with each other and other cells of the body plays an important role in regulating immune and inflammatory responses [Springer, T A. Nature, 346, 425, (1990); Springer, T. A. Cell 76, 301, (1994)]. Many of these interactions are mediated by specific cell surface molecules collectively referred to as cell adhesion molecules.

15 The adhesion molecules have been sub-divided into different groups on the basis of their structure. One family of adhesion molecules which is believed to play a particularly important role in regulating immune and inflammatory responses is the integrin family. This family of cell surface glycoproteins has a typical non-covalently linked heterodimer structure. At
20 least 14 different integrin alpha chains and 8 different integrin beta chains have been identified [Sonnenberg, A. Current Topics in Microbiology and Immunology, 184, 7, (1993)]. The members of the family are typically named according to their heterodimer composition although trivial nomenclature is widespread in this field. Thus the integrin termed $\alpha 4\beta 1$
25 consists of the integrin alpha 4 chain associated with the integrin beta 1 chain, but is also widely referred to as Very Late Antigen 4 or VLA4. Not all of the potential pairings of integrin alpha and beta chains have yet been observed in nature and the integrin family has been subdivided into a number of subgroups based on the pairings that have been recognised
30 [Sonnenberg, A. *ibid*].

The importance of cell adhesion molecules in human leukocyte function has been further highlighted by a genetic deficiency disease called Leukocyte Adhesion Deficiency (LAD) in which one of the families of
35 leukocyte integrins is not expressed [Marlin, S. D. *et al* J. Exp. Med. 164, 855 (1986)]. Patients with this disease have a reduced ability to recruit

leukocytes to inflammatory sites and suffer recurrent infections which in extreme cases may be fatal.

5 The potential to modify adhesion molecule function in such a way as to
beneficially modulate immune and inflammatory responses has been
extensively investigated in animal models using specific monoclonal
antibodies that block various functions of these molecules [e.g. Issekutz,
T. B. J. Immunol. 3394, (1992); Li, Z. *et al* Am. J. Physiol. 263, L723,
(1992); Binns, R. M. *et al* J. Immunol. 157, 4094, (1996)]. A number of
10 monoclonal antibodies which block adhesion molecule function are
currently being investigated for their therapeutic potential in human
disease.

One particular integrin subgroup of interest involves the $\alpha 4$ chain which
15 can pair with two different beta chains $\beta 1$ and $\beta 7$ [Sonnenberg, A. *ibid*].
The $\alpha 4\beta 1$ pairing occurs on many circulating leukocytes (for example
lymphocytes, monocytes and eosinophils) although it is absent or only
present at low levels on circulating neutrophils. $\alpha 4\beta 1$ binds to an adhesion
molecule (Vascular Cell Adhesion Molecule-1 also known as VCAM-1)
20 frequently up-regulated on endothelial cells at sites of inflammation
[Osborne, L. Cell, 62, 3, (1990)]. The molecule has also been shown to
bind to at least three sites in the matrix molecule fibronectin [Humphries,
M. J. *et al*. Ciba Foundation Symposium, 189, 177, (1995)]. Based on
data obtained with monoclonal antibodies in animal models it is believed
25 that the interaction between $\alpha 4\beta 1$ and ligands on other cells and the
extracellular matrix plays an important role in leukocyte migration and
activation [Yednock, T. A. *et al*, Nature, 356, 63, (1992); Podolsky, D. K.
et al. J. Clin. Invest. 92, 373, (1993); Abraham, W. M. *et al*. J. Clin.
Invest. 93, 776, (1994)].

30 The integrin generated by the pairing of $\alpha 4$ and $\beta 7$ has been termed
LPAM-1 [Holzmann, B and Weissman, I. EMBO J. 8, 1735, (1989)] and
like $\alpha 4\beta 1$, binds to VCAM-1 and fibronectin. In addition, $\alpha 4\beta 7$ binds to an
adhesion molecule believed to be involved in the homing of leukocytes to
35 mucosal tissue termed MAdCAM-1 [Berlin, C. *et al*, Cell, 74, 185, (1993)].
The interaction between $\alpha 4\beta 7$ and MAdCAM-1 may also be important at


sites of inflammation outside of mucosal tissue [Yang, X-D. *et al*, PNAS, 91, 12604 (1994)].

Regions of the peptide sequence recognised by $\alpha 4\beta 1$ and $\alpha 4\beta 7$ when they
5 bind to their ligands have been identified. $\alpha 4\beta 1$ seems to recognise LDV,
IDA or REDV peptide sequences in fibronectin and a QIDSP sequence in
VCAM-1 [Humphries, M. J. *et al*, *ibid*] whilst $\alpha 4\beta 7$ recognises a LDT
sequence in MAdCAM-1 [Briskin, M. J. *et al*, J. Immunol. 156, 719,
10 (1996)]. There have been several reports of inhibitors of these
interactions being designed from modifications of these short peptide
sequences [Cardarelli, P. M. *et al* J. Biol. Chem. 269, 18668, (1994);
Shroff, H. N. Bioorganic. Med. Chem. Lett. 6, 2495, (1996); Vanderslice,
P. J. Immunol. 158, 1710, (1997)]. It has also been reported that a short
15 peptide sequence derived from the $\alpha 4\beta 1$ binding site in fibronectin can
inhibit a contact hypersensitivity reaction in a trinitrochlorobenzene
sensitised mouse [Ferguson, T. A. *et al*, PNAS 88, 8072, (1991)].

Since the alpha 4 subgroup of integrins are predominantly expressed on
leukocytes their inhibition can be expected to be beneficial in a number of
20 immune or inflammatory disease states. However, because of the
ubiquitous distribution and wide range of functions performed by other
members of the integrin family it is very important to be able to identify
selective inhibitors of the alpha 4 subgroup.

25 We have now found a group of compounds which are potent and selective
inhibitors of $\alpha 4$ integrins. Members of the group are able to inhibit $\alpha 4$
integrins such as $\alpha 4\beta 1$ and/or $\alpha 4\beta 7$ at concentrations at which they
generally have no or minimal inhibitory action on α integrins of other
subgroups. The compounds are thus of use in medicine, for example in
30 the prophylaxis and treatment of immune or inflammatory disorders as
described hereinafter.

Thus according to one aspect of the invention we provide a compound of
formula (1)



()

wherein

- 5 R is (1) a group R¹X¹- where R¹ is an optionally substituted alkyl or aromatic group, and X¹ is a covalent bond or a -(CH₂)_n- [where n is an integer 1 or 2], -C(O)-, -CH₂C(O)-, -NHC(O)- or -SO₂- group, or (2) a group (Hal¹)₃CSO₂-, where Hal¹ is a fluorine or chlorine atom;
- R² and R³, which may be the same or different, is each a hydrogen or halogen atom or an alkyl, alkoxy, hydroxyl or nitro group;
- 10 Alk is an alkylene chain;
- m is zero or an integer 1;
- R⁴ is a hydrogen atom or a methyl group;
- R⁵ is a group -(CH₂)_pCO₂R⁸ where p is zero or an integer 1 and R⁸ is a hydrogen atom or an alkyl group;
- 15 R⁶ is a hydrogen atom or an alkyl group;
- Y is a sulphur atom or a -S(O)_q- group where q is an integer 1 or 2;
- X² is a -C(O)-, -C(O)O-, -CONH- or -S(O)₂- group;
- R⁷ is an optionally substituted alkyl group or an aryl or aralkyl group;
- and the salts, solvates and hydrates thereof.

20

It will be appreciated that compounds of formula (1) may have one or more chiral centres. Where one or more chiral centres is present, enantiomers or diastereomers may exist, and the invention is to be understood to extend to all such enantiomers, diastereomers and mixtures thereof, including racemates. Formula (1) and the formulae hereinafter are intended to represent all individual isomers and mixtures thereof, unless stated or shown otherwise.

25

30 In the compounds of formula (1), when the group R¹ is an optionally substituted alkyl group it may be for example an optionally substituted straight or branched chain C₁₋₆alkyl group such as an optionally substituted methyl or ethyl group. Optional substituents which may be

present on such groups include one, two or three halogen atoms, e.g. fluorine, chlorine, bromine or iodine atoms, or hydroxyl or C₁₋₄alkoxy e.g. methoxy or ethoxy groups.

- 5 Optionally substituted aromatic groups represented by the group R¹ in compounds of formula (1) include for example optionally substituted monocyclic or bicyclic fused ring C₆₋₁₂ aromatic groups, such as optionally substituted phenyl, 1- or 2-naphthyl, 1- or 2-tetrahydronaphthyl, indanyl or indenyl groups .

10

Optional substituents which may be present on aromatic groups of this type include one, two, three or more substituents selected from halogen atoms, e.g. fluorine, chlorine, bromine or iodine atoms, or C₁₋₆alkyl, e.g. methyl or ethyl, C₁₋₆alkoxy, e.g. methoxy or ethoxy, hydroxyl or nitro groups.

15

Alkyl groups represented by the groups R², R³, R⁶ and/or ,when present, R⁸ in compounds of the invention include for example straight or branched C₁₋₆alkyl groups such as methyl, ethyl, n-propyl, i-propyl, n-butyl, i-butyl, s-butyl or t-butyl groups.

20

Alkoxy groups represented by the groups R² and/or R³ include straight or branched C₁₋₆alkoxy groups such as methoxy or ethoxy groups.

25

The alkylene chain represented by Alk in compounds of formula (1) may be for example a straight or branched C₁₋₃alkylene chain such as a methylene or ethylene chain.

30

Optionally substituted alkyl groups represented by the group R⁷ in compounds of the invention include optionally substituted straight or branched C₁₋₆alkyl groups such as optionally substituted methyl, ethyl, n-propyl, i-propyl, n-butyl, i-butyl, s-butyl or t-butyl groups. Optional substituents which may be present on these groups include one, two or three substituents selected from halogen atoms, e.g. fluorine, chlorine, bromine or iodine atoms, or hydroxyl or C₁₋₄alkoxy, e.g. methoxy or ethoxy groups.

35

When in the compounds of the invention the group R^7 is an aryl group it may be for example an optionally substituted monocyclic or bicyclic fused ring C_{6-12} aromatic group as described above for the group R^1 .

- 5 When in the compounds of formula (1) R^7 is an aralkyl group it may be for example an optionally substituted monocyclic or bicyclic fused ring C_{6-12} aromatic C_{1-3} alkylene group. In groups of the type the aromatic portion may in particular be an optionally substituted aromatic group as described above for the group R^1 . The C_{1-3} alkylene portion may be for example a
 10 methylene or ethylene chain. Particular examples of aralkyl groups include optionally substituted benzyl groups.

The presence of certain substituents in the compounds of formula (1) may enable salts of the compounds to be formed. Suitable salts include
 15 pharmaceutically acceptable salts, for example salts derived from inorganic and organic bases. Particular examples of such salts include alkali metal salts such as sodium or potassium salts, alkaline earth metal salts such as magnesium or calcium salts, and organic amine salts such as morpholine, piperidine, dimethylamine or diethylamine salts.

20 One particular class of compounds according to the invention has the formula (1) wherein R^4 is a hydrogen atom and the remaining groups are as defined above for formula (1) and the salts, solvates and hydrates thereof.

25 In another class of compounds of formula (1) the group R^5 is a $-CH_2CO_2H$ group, or in particular is a $-CO_2H$ group.

30 In compounds of this class, and in general in compounds of formula (1) Y is preferably a sulphur atom.

In another general preference, m in compounds of formula (1) is the integer 1 and Alk in particular is a $-CH_2-$ group. In compounds of this type, and when R^5 is a $-CO_2H$ group, the carbon atom to which R^5 and Alk are
 35 attached forms a chiral centre and is preferably in the L configuration.

The group R in compounds of formula (1) is preferably a R^1X^1 - group. In compounds of this type R^1 is preferably an optionally substituted phenyl group. Particularly useful groups of this type include mono-, di- or trisubstituted phenyl groups. The substituent(s) may be located on any available carbon atom in the phenyl ring at the 2-, 3-, 4-, 5- and 6-positions relative to the point of attachment of the phenyl group to the remainder of the molecule of formula (1). Thus, for example when more than one substituent is present, the substituents may be located at the 2,6- and 2, 4, 6- positions. Particularly useful substituents include halogen atoms such as chlorine and fluorine atoms. X^1 in compounds of these particular types is preferably a $-CH_2-$ or $-C(O)-$ group.

R^6 in compounds of formula (1) may for example be a methyl group or in particular a hydrogen atom.

X^2 in the compounds according to the invention is preferably a $-C(O)-$ group.

The group R^7 in the compounds according to the invention is preferably a methyl group.

Particularly useful compounds according to the invention include:
N-Acetyl-*D*-thioprolin-(*O*-2,6-dichlorobenzyl)-*L*-tyrosine;
N-Acetyl-*D*-thioprolin-(*O*-2,4,6-trichlorobenzyl)-*L*-tyrosine;
N-Acetyl-*D*-thioprolin-(*O*-2,6-difluorobenzyl)-*L*-tyrosine;
and the salts, solvates and hydrates thereof.

Compounds according to the invention are potent and selective inhibitors of α_4 integrins. The ability of the compounds to act in this way may be simply determined by employing tests such as those described in the Examples hereinafter.

The compounds are of use in modulating cell adhesion and in particular are of use in the prophylaxis and treatment of diseases or disorders involving inflammation in which the extravasation of leukocytes plays a role. Diseases or disorders of this type include inflammatory arthritis such

as rheumatoid arthritis vasculitis or polydermatomyositis, multiple sclerosis, allograft rejection, diabetes, inflammatory dermatoses such as psoriasis or dermatitis, asthma and inflammatory bowel disease. The compounds may also be useful for modulating the circulating levels of
5 early haematopoietic cells, such as stem cells to enable their collection for e.g. bone marrow transplantation.

For the prophylaxis or treatment of disease the compounds according to the invention may be administered as pharmaceutical compositions, and
10 according to a further aspect of the invention we provide a pharmaceutical composition which comprises a compound of formula (1) together with one or more pharmaceutically acceptable carriers, excipients or diluents.

Pharmaceutical compositions according to the invention may take a form
15 suitable for oral, buccal, parenteral, nasal, topical or rectal administration, or a form suitable for administration by inhalation or insufflation.

For oral administration, the pharmaceutical compositions may take the form of, for example, tablets, lozenges or capsules prepared by
20 conventional means with pharmaceutically acceptable excipients such as binding agents (e.g. pregelatinised maize starch, polyvinylpyrrolidone or hydroxypropyl methylcellulose); fillers (e.g. lactose, microcrystalline cellulose or calcium hydrogen phosphate); lubricants (e.g. magnesium stearate, talc or silica); disintegrants (e.g. potato starch or sodium
25 glycollate); or wetting agents (e.g. sodium lauryl sulphate). The tablets may be coated by methods well known in the art. Liquid preparations for oral administration may take the form of, for example, solutions, syrups or suspensions, or they may be presented as a dry product for constitution with water or other suitable vehicle before use. Such liquid preparations
30 may be prepared by conventional means with pharmaceutically acceptable additives such as suspending agents, emulsifying agents, non-aqueous vehicles and preservatives. The preparations may also contain buffer salts, flavouring, colouring and sweetening agents as appropriate.

35 Preparations for oral administration may be suitably formulated to give controlled release of the active compound.

For buccal administration the compositions may take the form of tablets or lozenges formulated in conventional manner.

- 5 The compounds for formula (1) may be formulated for parenteral administration by injection e.g. by bolus injection or infusion. Formulations for injection may be presented in unit dosage form, e.g. in glass ampoule or multi dose containers, e.g. glass vials. The compositions for injection may take such forms as suspensions, solutions or emulsions in oily or aqueous vehicles, and may contain formulatory agents such as suspending, stabilising, preserving and/or dispersing agents. Alternatively, the active ingredient may be in powder form for constitution with a suitable vehicle, e.g. sterile pyrogen-free water, before use.
- 10
- 15 In addition to the formulations described above, the compounds of formula (1) may also be formulated as a depot preparation. Such long acting formulations may be administered by implantation or by intramuscular injection.
- 20 For nasal administration or administration by inhalation, the compounds for use according to the present invention are conveniently delivered in the form of an aerosol spray presentation for pressurised packs or a nebuliser, with the use of suitable propellant, e.g. dichlorodifluoromethane, trichlorofluoromethane, dichlorotetrafluoroethane, carbon dioxide or other suitable gas or mixture of gases.
- 25

The compositions may, if desired, be presented in a pack or dispenser device which may contain one or more unit dosage forms containing the active ingredient. The pack or dispensing device may be accompanied by instructions for administration.

30

The quantity of a compound of the invention required for the prophylaxis or treatment of a particular condition will vary depending on the compound chosen, and the condition of the patient to be treated. In general, however, daily dosages may range from around 100ng/kg to 100mg/kg e.g. around 0.01mg/kg to 40mg/kg body weight for oral or buccal

35

administration, from around 10ng/kg to 50mg/kg body weight for parenteral administration and around 0.05mg to around 1000mg e.g. around 0.5mg to around 1000mg for nasal administration or administration by inhalation or insufflation.


5

The compounds of the invention may be prepared by a number of processes as generally described below and more specifically in the Examples hereinafter. In the following process description, the symbols R, R¹-R⁷, Alk, m, Y and X¹ when used in the formulae depicted are to be understood to represent those groups described above in relation to formula (1) unless otherwise indicated. In the reactions described below, it may be necessary to protect reactive functional groups, for example hydroxy or carboxy groups, where these are desired in the final product, to avoid their unwanted participation in the reactions. Conventional protecting groups may be used in accordance with standard practice [see, for example, Green, T. W. in "Protective Groups in Organic Synthesis", John Wiley and Sons, 1991]. In some instances, deprotection may be the final step in the synthesis of a compound of formula (1) and the processes according to the invention described hereinafter are to be understood to extend to such removal of protecting groups.

15
20

Thus according to a further aspect of the invention, a compound of formula (1) wherein R⁵ is a group $-(CH_2)_pCO_2R^8$ in which p is zero or an integer 1 and R⁸ is an alkyl group may be prepared by coupling an amine of formula (2):


25



(2)

(where R⁵ is as just described) or a salt thereof with an acid of formula (3):

30



(3)

or an active derivative thereof.

- 5 Active derivatives of acids of formula (3) include anhydrides, esters and halides. Particular esters include pentafluorophenyl or succinyl esters.


10 The coupling reaction may be performed using standard conditions for reactions of this type. Thus for example the reaction may be carried out in a solvent, for example an inert organic solvent such as an amide, e.g. a substituted amide such as dimethylformamide, an ether, e.g. a cyclic ether such as tetrahydrofuran, or a halogenated hydrocarbon, such as dichloromethane, at a low temperature, e.g. around $-30^\circ C$ to around ambient temperature, optionally in the presence of a base, e.g. an organic base such as an amine, e.g. triethylamine, pyridine, or a cyclic amine, such as N-methylmorpholine.

20 Where an acid of formula (3) is used, the reaction may additionally be performed in the presence of a condensing agent, for example a diimide such as 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide or N,N'-dicyclohexylcarbodiimide, advantageously in the presence of a catalyst such as a N-hydroxy compound e.g. a N-hydroxytriazole such as 1-hydroxybenzotriazole. Alternatively, the acid may be reacted with a chloroformate, for example ethylchloroformate, prior to reaction with the amine of formula (2).

30 Intermediate amines of formula (2) wherein R^5 is a group $-CO_2R^8$ are either known compounds or may be prepared from known starting materials by methods analogous to those used for the preparation of the known compounds. Where appropriate, standard substitution approaches such as those described below employing for example alkylation, arylation, acylation, halogenation, sulfonylation, nitration or coupling

reactions may be used to obtain new R, R², R³ or R⁵ substituents in known amines of formula (2). In these reactions the amine may need to be suitably protected, for example as described by Green, T. W. *ibid*.

- 5 Intermediate amines of formula (2) wherein m is zero and R⁵ is a group -CH₂CO₂R⁸ may be prepared by reaction of an aldehyde or ketone of formula (4):




(4)

10

with malonic acid and ammonia or an ammonium salt, e.g. ammonium acetate, in the presence of a base, followed by reaction with an alcohol R⁸OH in the presence of an acid such as hydrochloric acid.

- 15 Intermediate amines of formula (2) wherein R⁵ is a group -CH₂CO₂R⁸ may be prepared by reaction of an α,β -unsaturated ester of formula (5):




(5)

- 20 with benzylamine or a substituted benzylamine, optionally in the presence of a base, followed by hydrogenation using for example hydrogen or a hydrogen donor such as formic acid and a transfer agent such as a metal catalyst, for example palladium on a support such as carbon in a solvent such as methanol at an ambient or elevated temperature.


25

The esters of formula (5) may be obtained by reaction of an aldehyde or ketone of formula (6):



with a phosphonium salt $(\text{R}^9)_3\text{P}^+\text{CH}_2\text{CO}_2\text{R}^8\text{Hal}^-$ (where Hal is a halogen atom and R^9 is for example a phenyl group) or a stabilised ylide $(\text{R}^9)_3\text{P}=\text{CHCO}_2\text{R}^8$ in the presence of a base such as sodium ethoxide in a solvent such as ethanol or phenyllithium in a solvent such as tetrahydrofuran at around ambient temperature.

Aldehydes of formula (6) wherein R^4 is a hydrogen atom may be prepared by reduction of a corresponding ester of formula (7):



(where R^8 is an alkyl group) using a reducing agent such as a metal hydride, e.g. diisobutylaluminium hydride, at a low temperature e.g. around -78°C in an organic solvent such as toluene.


Ketones of formula (6) wherein R^4 is a methyl group may be prepared by treating a corresponding aldehyde of formula (6) with a Grignard reagent such as methylmagnesium bromide, or methyllithium in a solvent such as tetrahydrofuran, at a low temperature e.g. around -55°C to 0°C and oxidising the resulting alcohol using an oxidising agent such as manganese dioxide.

Intermediate aldehydes and ketones of formula (4) and intermediate esters of formula (7) are either known compounds or may be prepared from known starting materials by methods analogous to those used for the preparation of the known compounds, where appropriate employing

standard substitution approaches to obtain any desired R , R^2 , R^3 and/or R^8 group as described above in relation to the intermediate amines of formula (2).

- 5 The acids of formula (3) for use in the preparation of compounds of the invention are also either known compounds or may be prepared from known starting materials by use of analogous processes to those used for the preparation of the known compounds, for example by acylation or sulphonylation of an acid of formula (8):

10




- or a protected derivative thereof using for example a reagent R^7COHal , R^7CO_2H , R^7SO_2Hal or R^7NCO and standard conditions for reactions of this type such as those described hereinafter for the functionalisation of phenols of formula (9).

- 15 In another aspect of the invention a compound of formula (1) may be obtained from a corresponding compound of formula (1) via an inter-conversion process.

- 20 Thus, in one particular example, a compound of formula (1) wherein R^5 is a $-CO_2H$ or $-CH_2CO_2H$ group may be obtained by hydrolysis of a corresponding ester wherein R^5 is a $-CO_2R^8$ or $-CH_2CO_2R^8$ group and R^8 is an alkyl group. The hydrolysis may be performed using either an acid or a base depending on the nature of the ester starting material, for example an organic acid such as trifluoroacetic acid or an inorganic base such as lithium hydroxide optionally in an aqueous organic solvent such as an amide, e.g. a substituted amide such as dimethylformamide, an ether, e.g. a cyclic ether such as tetrahydrofuran or dioxane or an alcohol, e.g. methanol at around ambient temperature. Where desired, mixtures of such solvents may be used.

In a still further aspect of the invention a compound of formula (1) may also be prepared by functionalisation of a phenol of formula (9):



5 using standard substitution approaches employing for example alkylation, arylation, acylation, sulphonylation or coupling reactions. In these reactions the starting materials of formula (9) may first be obtained by use of the appropriate phenol intermediates in the reactions previously
10 described to obtain compounds of formula (1).

Thus in one example, a phenol of formula (9) may be alkylated or arylated using a reagent R^1X^1L in which X^1 is a covalent bond or a $-(CH_2)_n$ group and L is a leaving atom or group such as a halogen atom, e.g. a fluorine, bromine, iodine or chlorine atom or a sulphonyloxy group such as an
15 alkylsulphonyloxy, e.g. trifluoromethylsulphonyloxy or arylsulphonyloxy, e.g. p-toluenesulphonyloxy group.

The alkylation or arylation reaction may be carried out in the presence of a
20 base such as a carbonate, e.g. caesium or potassium carbonate, an alkoxide, e.g. potassium t-butoxide, or a hydride, e.g. sodium hydride, in a dipolar aprotic solvent such as an amide, e.g. a substituted amide such as dimethylformamide or an ether, e.g. a cyclic ether such as tetrahydrofuran at for example around 0°C to around ambient temperature. The
25 compounds R^1X^1L are either known and readily available or may be obtained by simple manipulation of known compounds, for example as described in the Examples hereinafter.


In another example, a phenol of formula (9) may be functionalised by
30 acylation, for example by reaction with a reagent $R^1X^1\text{Hal-}$ [wherein X^1 is a $-C(O)-$, $-CH_2C(O)-$ or $-NHC(O)-$ group and Hal is a halogen atom such as a chlorine atom] in the presence of a base, such as a tertiary amine,

e.g. triethylamine, in a solvent such as a halogenated hydrocarbon, e.g. dichloromethane or carbon tetrachloride, at for example ambient temperature.

- 5 In a further example a compound of the invention may be obtained by sulphonylation of a phenol of formula (9) by reaction with a reagent R^1X^1L or $(Hal^1)_3CSO_2L$ [in which X^1 is $-SO_2-$ and L is a leaving group as defined above] in the presence of a base, for example an inorganic base such as sodium hydride in a solvent such as an amide, e.g. a substituted amide
10 such as dimethylformamide at for example ambient temperature.

- In another example a compound of formula (1) wherein R is a group R^1X^1 [where X^1 is a covalent bond or a $-(CH_2)_n$ group] may be obtained by coupling a phenol of formula (9) with a reagent R^1OH or $R^1(CH_2)_nOH$ in a
15 solvent such as tetrahydrofuran in the presence of a phosphine, e.g. triphenylphosphine and an activator such as diethyl-, diisopropyl- or dimethylazodicarboxylate.

- In a further process according to the invention a compound of formula (1)
20 may be prepared by acylation or sulphonylation of an intermediate of formula (10):



- 25 Reagents for these reactions include for example compounds of the types R^7COHal , R^7CO_2H , R^7SO_2Hal or R^7NCO . The reactions may be performed using standard conditions such as those described above in relation to the acylation or sulphonylation of phenols of formula (9). It will be appreciated that in some instances and under suitable conditions the
30 reaction may also be performed on compounds of formula (10) in which R is a hydrogen atom so that acylation or sulphonylation takes place at both ends of the molecule. In general in this process any carboxyl group in intermediates of formula (10) will need to be protected, for example as a

methyl ester, and, where required, the free acid subsequently regenerated by hydrolysis as described herein.

5 Salts of compounds of formula (1) may be prepared by reaction of a compound of formula (1) with an appropriate base in a suitable solvent or mixture of solvents e.g. an organic solvent such as an ether e.g. diethylether, or an alcohol, e.g. ethanol using conventional procedures.

10 Where it is desired to obtain a particular enantiomer of a compound of formula (1) this may be produced from a corresponding mixture of enantiomers using any suitable conventional procedure for resolving enantiomers.

15 Thus for example diastereomeric derivatives, e.g. salts, may be produced by reaction of a mixture of enantiomers of formula (1) e.g. a racemate, and an appropriate chiral compound, e.g. a chiral base. The diastereomers may then be separated by any convenient means, for example by crystallisation and the desired enantiomer recovered, e.g. by treatment with an acid in the instance where the diastereomer is a salt.

20 In another resolution process a racemate of formula (1) may be separated using chiral High Performance Liquid Chromatography. Alternatively, if desired a particular enantiomer may be obtained by using an appropriate chiral intermediate in one of the processes described above.

25 The following Examples illustrate the invention. All temperatures are in °C. The following abbreviations are used:

30	EDC - 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide;	
	DMF - dimethylformamide;	DMSO - dimethylsulphoxide;
	HOBT - 1-hydroxybenzotriazole;	THF - tetrahydrofuran;
	TFA - trifluoroacetic acid;	NMM - N-methylmorpholine;
	DCM - dichloromethane;	Ph - phenyl;
	tyr - tyrosine;	Ar - aryl;
35	thiopro - thioproline;	pyr - pyridine

INTERMEDIATE 1**N-Acetyl-D-thiopropine-L-tyrosine tert.butyl ester**

EDC (4.22g, 22mmol) was added to a solution of N-acetyl-D-thiopropine (3.50g, 20mmol), tyrosine tert.butyl ester (4.74g, 20mmol), HOBT (2.97g, 22mmol) and NMM (2.42ml, 22mmol) in DMF (80ml) at 0°. The mixture was stirred at room temperature overnight. The DMF was evaporated *in vacuo* and the residue dissolved in ethyl acetate (600ml) and water (50ml). The organic phase was washed with 10% citric acid (150ml), saturated aqueous NaHCO₃ (150ml), water (150ml) and brine (150ml), dried (Na₂SO₄) and evaporated *in vacuo* to give the title compound as an off-white solid (7.39g, 94%); δ H (DMSO-d₆, 300K) (2 rotameric species observed) 9.17 (1H, br s, ArCH), 8.43 (3, \downarrow 8.0Hz), and 8.09 (d, \downarrow 8.0Hz) together (1H, CONH), 6.98 (2H, t, \downarrow 7.6, ArH), 6.64 (2H, d, \downarrow 7.3, ArH), 4.81-4.67 (m), 4.47 (d, \downarrow 8.7Hz) and 4.35-4.21 (m), together (4H, CH α -tyr, CH α thioprop and NCH₂S), 3.32 (dd, \downarrow 7.2, 11.6Hz), 3.17 (dd, \downarrow 7.5, 11.6Hz), 2.99-2.72 (m) together (4H, CH₂Ar + CHCH₂S), 2.06 (s) and 1.84 (s) together (3H, CH₃CO) and 1.36 (9H, s, CO₂^tBu); m/z (ESI, 15V) 395 ($M^+ + 1$).

INTERMEDIATE 2**N-Acetyl-D-thiopropine-L-tyrosine methyl ester**

EDC (2.11g, 11mmol) was added to a stirred solution of N-acetyl-D-thiopropine (1.75g, 10mmol), tyrosine methyl ester (2.32g, 10mmol), HOBT (1.49g, 11mmol) and NMM (2.31ml, 21mmol) in DMF (50ml) at 0°. The mixture was stirred at room temperature overnight. The DMF was evaporated *in vacuo* and the residue dissolved in ethyl acetate (400ml), and water (50ml). The organic phase was washed with 10% citric acid (100ml), saturated aqueous NaHCO₃ (100ml), water (100ml) and brine (100ml), dried (Na₂SO₄) and evaporated *in vacuo* to give the title compound as a slightly yellow powdery solid (2.75g, 78%), used without further purification. A small portion was crystallised from EtOAc to give a white microcrystalline solid, m.p. 189-190°. δ H (DMSO-d₆, 400K) 7.7 (1H, br s, CONH), 6.98 (2H, d, \downarrow 8.6Hz, ArH), 6.69 (2H, d, \downarrow 8.5Hz, ArH), 4.83 (1H, dd, \downarrow 3.9, 7.4Hz, CH α thioprop), 4.77 (1H, d, \downarrow 9.2 NCH_AH_BS), 4.53 (1H, dt, \downarrow 5.8, 8.1Hz, CH α tyr), 4.38 (1H, d, \downarrow 9.2Hz, NCH_AH_BS), 3.65 (3H, s, C O₂HC), 3.25 (1H, dd, \downarrow 7.3, 11.5Hz, CHCH_AH_BS), 3.05-2.97 (2H, m,

$\text{CH}_A\text{H}_B\text{Ar} + \text{CHCH}_A\text{H}_B\text{S}$), 2.89 (1H, dd, J 8.2, 14.1Hz, $\text{CH}_A\text{H}_B\text{Ar}$) and 1.99 (3H, s, CH_3CO) [phenolic OH not observed at 400K, for δH (DMSO-d_6 , 300K) 9.19 (1H, br s, ArOH)]; m/z ESI, 27V), 353 ($\text{M}^+ + 1$).

5 INTERMEDIATE 3

3-Nitro-L-tyrosine methyl ester hydrochloride

Acetyl chloride (8.9ml, 125mmol) was added slowly to methanol (100ml) at 0° . 3-Nitro-L-tyrosine (5.65g, 25mmol) was added and the mixture refluxed for 2h. The solvent was evaporated *in vacuo* and the yellow solid obtained recrystallised from methanol to give the title compound as yellow needles (2.97g, 43%), m.p. 200-201 $^\circ$. (Found: C, 43.29; H, 4.69; N, 10.19. $\text{C}_{10}\text{H}_{12}\text{N}_2\text{O}_5 \cdot \text{HCl}$ requires C, 43.41; H, 4.74; N, 10.13%); δH (CD_3OD) 8.00 (1H, d, J 2.2Hz, ArH), 7.53 (1H, dd, J 2.3, 8.6Hz, ArH), 7.17 (1H, d, J 8.6Hz, ArH), 4.37 (1H, dd, J 6.4, 7.0Hz, CH_α), 3.83 (3H, s, CO_2CH_3), 3.29 (1H, dd, J 6.3, 14.6 Hz, $\text{CH}_A\text{H}_B\text{Ar}$) and 3.20 (1H, dd, J 7.1, 14.6 Hz, $\text{CH}_A\text{H}_B\text{Ar}$); m/z (ES, 27V) 241 ($\text{M}^+ + 1$).

INTERMEDIATE 4

N-Acetyl-D-thiopropine-3-nitro-L-tyrosine methyl ester

EDC.HCl (1.056g, 5.5mmol) was added to a solution of *N*-acetyl-D-thiopropine (875mg, 5mmol), HOBT (743mg, 5.5mmol) and NMM (1.15ml, 10.5mmol) in DMF (25ml) at room temperature. The reaction mixture was stirred overnight. The DMF was evaporated *in vacuo* and the residue dissolved in ethyl acetate (150ml) and water (50ml). The organic phase was washed with hydrochloric acid (1M, 30ml), saturated aqueous NaHCO_3 (30ml), water (50ml) and brine (30ml), dried (Na_2SO_4) and evaporated *in vacuo* to give the title compound as a yellow foam (1.59g, 80%); δH (DMSO-d_6 , 300K) (2 rotameric species observed) 10.79 (s) and 10.76 (s) together (1H, ArOH), 8.61 (d, J 8.2Hz) and 8.32 (d, J 8.3Hz) together (1H, CONH), 7.75-7.71 (1H, m, ArH), 7.44-7.37 (1H, m, ArH), 7.06-7.01 (1H, m, ArH), 4.74-4.67 (m) and 4.56-4.43 (m) and 4.44 (d, J 8.6Hz), and 4.19 (d, J 9.7Hz) together (4H, $2 \times \text{CH}_\alpha + \text{NCH}_2\text{S}$), 3.64 (3H, s, CO_2CH_3), 3.33-2.77 (4H, m, $\text{CHCH}_2\text{S} + \text{CH}_2\text{Ar}$) and 2.03 (s) and 1.84 (s) together (3H, CH_3CO); m/z (ES, 30V) 398 ($\text{M}^+ + 1$).

INTERMEDIATE 5 **α -Methyl-*L*-tyrosine methyl ester hydrochloride**

Anhydrous hydrogen chloride was bubbled through a solution of α -methyl-*L*-tyrosine (1g, 5.13mmol) in methanol (100ml) for a few minutes and the solution stirred at room temperature for 48h. The solvent was evaporated *in vacuo* and the residue freeze dried from a mixture of methanol and water to give the title compound as a white powder (1.28g, 100%); δ H (CD₃OD) 7.00 (2H, d, J 8.6Hz, ArH), 6.78 (2H, d, J 8.6Hz, ArH), 3.82 (3H, s, CO₂CH₃), 3.19 (1H, d, J 14.3Hz, CH_AH_BAr), 3.01 (1H, d, J 14.3Hz, CH_AH_BAr) and 1.59 (3H, s, CCH₃); m/z (ES, 27V) 210 ($M^+ + 1$).

INTERMEDIATE 6***N*-Acetyl-*D*-thiopropine- α -methyl-*L*-tyrosine methyl ester**

EDC.HCl (760mg, 3.96mmol) was added to a solution of *N*-acetyl-*D*-thiopropine (630mg, 3.6mmol), Intermediate 5 (880mg, 3.6mmol), HOBT (535mg, 3.96mmol) and NMM (834 μ l, 7.6mmol) in DMF (20ml). The mixture was stirred at room temperature overnight. The DMF was evaporated *in vacuo* and the residue dissolved in ethyl acetate (150ml) and water (50ml). The organic phase was washed with 10% citric acid (50ml), saturated aqueous NaHCO₃ (50ml) and water (50ml), dried (Na₂SO₄) and evaporated *in vacuo*. The residue was purified by column chromatography (SiO₂, methanol/DCM, 7:93) to give the title compound as a colourless gum (680mg, 52%); δ H (DMSO-*d*₆, 400K) 7.40 (1H, br s, CONH), 6.93 (2H, d, J 8.6Hz, ArH), 6.68 (2H, d, J 8.3Hz, ArH), 4.84 (1H, dd, J 3.8, 7.4Hz, CH α thiopro), 4.78 (1H, d, J 9.2Hz, NCH_AH_BS), 4.40 (1H, d, J 9.2Hz, NCH_AH_BS), 3.63 (3H, s, CO₂CH₃), 3.30 (1H, dd, J 7.4, 11.5Hz, CHCH_AH_BS), 3.16 (1H, dd, J 3.8, 11.5Hz, CHCH_AH_BS), 3.13 (1H, d, J 13.7Hz, CH_AH_BAr), 3.03 (1H, d, J 13.7Hz, CH_AH_BAr), 2.02 (3H, s, CH₃CO) and 1.38 (3H, s, CCH₃); m/z (ES, 15V) 367 ($M^+ + 1$).

INTERMEDIATE 7**2,4,6-Trichlorobenzyl alcohol**

A solution of lithium aluminium hydride (1M in THF, 18mmol, 18ml) was added slowly to a solution of 2,4,6-trichlorobenzoyl chloride (4.35g, 17.8mmol) in THF (70ml) at 0°. After 1h water (685 μ l) was added, followed by aqueous sodium hydroxide (3M, 685 μ l) and more water

(2.06ml). The suspension was stirred vigorously for 1h, the precipitate filtered off and the filtrate evaporated *in vacuo* to give a slightly yellow solid. Recrystallisation from diisopropylether gave the title compound as white needles (2.63g, 70%), m.p. 100-101°. δ H (CDCl₃) 7.35 (2H, s, ArH), 4.91 (2H, br s, CH₂CH) and 2.07 (1H, br s, CH); m/z (ES, 60V) 193 (M-OH).

INTERMEDIATE 8

2,4,6-Trichlorobenzyl bromide

Thionyl bromide (682 μ l, 8.8mmol) was added to a solution of Intermediate 7 (846mg, 4mmol) in DCM (20ml) at 0°. The reaction was stirred at room temperature overnight then quenched with water. The mixture was diluted with DCM (100ml), washed with saturated aqueous NaHCO₃ (30ml) and water (30ml), dried (Na₂SO₄) and evaporated *in vacuo*. The residue was purified by column chromatography (short plug SiO₂, hexane) to give the title compound as a colourless oil which crystallised on standing to white crystals (982mg, 89%), m.p. 51-52°. δ H (CDCl₃) 7.36 (2H, s, ArH) and 4.70 (2H, s, CH₂Br).

EXAMPLE 1

N-Acetyl-D-thiopropine-(O-2,6-dichlorobenzoyl)-L-tyrosine tert.butyl ester

A solution of Intermediate 1 (705mg, 1.79mmol) in THF (5ml) was added to a suspension of sodium hydride (60% in oil, 79mg, 1.97mmol) and 2,6-dichlorobenzoyl chloride (283 μ l, 1.97mmol) in THF (10ml) at 0°. The mixture was stirred at room temperature for 6h then quenched with aqueous NH₄Cl (5ml). The mixture was extracted with DCM (2 x 75ml) and the combined organic extracts dried (Na₂SO₄) and evaporated *in vacuo*. The residue was purified by chromatography (SiO₂; ethyl acetate/hexane 80:20) to give the title compound as a white foam (920mg, 91%); δ H (DMSO-d₆, 300K) (2 rotameric species observed) 8.55 (d, \downarrow 7.9Hz), and 8.25 (d, \downarrow 8.0Hz), together (1H, NHCO), 7.70-7.59 (3H, m, COArH), 7.38-7.34 (2H, m, CH₂ArH), 7.21-7.17 (2H, m, CHArH), 4.80-4.67 (m) and 4.50-4.34 (m) and 4.48 (d, \downarrow 8.7Hz) and 4.23 (d, \downarrow 9.6Hz) together (4H, 2 x CH α + NCH₂S), 3.30-2.80 (4H, m, CH₂Ar + CHCH₂S), 2.07 (s)

and 1.83 (s) together (3H, CH₃CO) and 1.38 (9H, s, CO₂tBu); m/z (ESI, 15V) 567 ($M^+ + 1$).

EXAMPLE 2

5 *N*-Acetyl-*D*-thiopropine-(*O*-2,6-dichlorobenzoyl)-*L*-tyrosine

A solution of the compound of Example 1 (910mg, 1.60mmol) in a mixture of TFA/H₂O (9:1, 20ml) was stirred at room temperature for 2h. The solvents were evaporated *in vacuo* and the residue freeze dried from a mixture of methanol/H₂O to give the title compound as a fluffy white solid
 10 (809mg, 99%) δ H (DMSO-*d*₆, 400K) 7.75 (1H, br d, CONH), 7.62-7.53 (3H, m, COArH), 7.34 (2H, d, \downarrow 8.7Hz, CH₂-ArH), 7.20 (2H, d, \downarrow 8.7Hz, CH₂ArH), 4.82 (1H, dd, \downarrow 3.9, 7.3Hz, CH α thioprop), 4.76 (1H, d, \downarrow 9.2Hz CH_AH_BS), 4.57 (1H, dt, \downarrow 5.4, 8.4Hz, CH α tyr), 4.37 (1H, d, \downarrow 9.1Hz, NCH_AH_BS), 3.25 (1H, dd, \downarrow 7.4, 11.6Hz, CHCH_AH_BS), 3.19 (1H, dd, \downarrow 5.3, 14.1Hz, CH_AH_BAr), 4.07-2.99 (2H, m, CHCH_AH_BS + CH_AH_BAr) and 1.98
 15 (3H, s, CH₃CO); m/z (ESI, 27V) 511 ($M^{++} + 1$); $[\alpha]_D^{24.5} = +76.53^\circ$ (c, = 0.69, methanol).

EXAMPLE 3

20 *N*-Acetyl-*D*-thiopropine - (*O*-2,6-dimethoxybenzoyl)-*L*-tyrosine methyl ester

A solution of Intermediate 2 (352mg, 1mmol) in DMF (5ml) was added to a suspension of sodium hydride (60% in oil, 44mg, 1.1mmol) in DMF (3ml) at 0°. After 5 min at room temperature a yellow solution was obtained, to
 25 this was added a solution of 2,6-dimethoxybenzoyl chloride (241mg, 1.22mmol) in DMF (2ml). The mixture was stirred for 1h then quenched with water and the DMF evaporated *in vacuo*. The residue was dissolved in ethyl acetate (100ml) and washed in water (3 x 30ml) and brine (30 ml), dried (Na₂SO₄) and evaporated *in vacuo*. The residue was purified by
 30 chromatography (SiO₂, DCM/methanol, 95:5) to give the title compound as a yellow gum (462mg, 90%); δ H (DMSO-*d*₆, 300K) (2 rotameric species observed) 8.63 (3, \downarrow 8.2Hz) and 8.38 (d, \downarrow 8/1Hz) together (1H, CONH), 7.43 (1H, t, \downarrow 8.4Hz, CO-Ar_fH), 7.29 (2H, t, \downarrow 8.1Hz, CH₂ArH), 7.09-7.06 (2H, m, CH₂ArH), 6.77 (2H, d, \downarrow 8.4Hz, CO-Ar_mH), 4.79-4.68 (m) and
 35 4.60-4.44 (m) and 4.45 (d, \downarrow 8.7Hz) and 4.23 (d, \downarrow 9.7Hz) together (4H, 2 x CH α + NCH₂S), 3.84 (6H, s, 2 x ArOMe), 3.65 (s) and 3.64 (s) together

(3H, CO₂Me), 3.32-2.73 (4H, m, CH₂Ar + CHCH₂S) and 2.06 (s) and 1.84 (s) together (3H, CH₃CO); m/z (ESI, 15V) 517 ($M^+ + 1$).

EXAMPLE 4

5 N-Acetyl-D-thiopropine - (O-2,6-dimethoxybenzoyl)-L-tyrosine

Lithium hydroxide (41mg, 0.97mmol) was added to a solution of the compound of Example 3 (455mg, 0.88mmol) in a mixture of THF (10ml) and water (10ml). The mixture was stirred at room temperature for 10 min then the THF was evaporated *in vacuo*. The residue was purified by
10 chromatography (SiO₂; DCM/methanol/acetic acid, 90:5:5). The gum obtained was freeze-dried from a mixture of methanol and water to give the title compound as a fluffy white solid (401mg, 91%). δ H (DMSO-d₆, 400K) 7.7 (1H, br d, CONH), 7.40 (1H, t, \downarrow 8.4Hz, COAr_fH), 7.28 (2H, d, \downarrow 8.7Hz, CH₂Ar_H), 7.09 (2H, d, \downarrow 8.6Hz, CH₂Ar_H), 6.77 (2H, d, \downarrow 7.8Hz, COAr_mH), 4.83 (1H, dd, \downarrow 4.0, 7.3 Hz, CH α thioprop), 4.77 (1H, d, \downarrow 9.2Hz, NCH_AH_BS), 4.57-4.51 (1H, m, CH α tyr), 4.38 (1H, d, \downarrow 98.2Hz, NCH_AH_BS), 3.87 (6H, s, 2 x COArOMe), 3.25 (1H, dd, \downarrow 7.4, 11.5Hz, CHCH_AH_BS), 3.17 (1H, dd, \downarrow 5.4, 14.1 CH_AH_BAr), 3.06-2.98 (2H, m, CHCH_AH_BS + CH_AH_BAr) and 1.99 (3H, s, CH₃CO), m/z (ESI, 15V) 503 ($M^+ + 1$).

20 EXAMPLE 5

N-Acetyl-D-thiopropine - (O-benzyl)-L-tyrosine methyl ester

EDC (211mg, 1.1mmol) was added to a stirred solution of N-acetyl-D-thiopropine (175mg, 1mmol), O-benzyl tyrosine methyl ester hydrochloride
25 (322mg, 1mmol), HOBT (149mg, 1.1mmol) and NMM (242 μ l, 2.2mmol) in DCM (10ml) at 0°. The mixture was stirred at room temperature overnight then diluted with DCM (100ml). The DCM solution was washed with 1M hydrochloric acid (30ml), saturated aqueous NaHCO₃ (30ml) and water (30ml), dried (Na₂SO₄) and evaporated *in vacuo*. The residue was
30 purified by chromatography (SiO₂; ethyl acetate) to give the title compound as a white foam (417mg, 94%); δ H (DMSO-d₆, 300K) (2 rotameric species observed) 8.55 (d, \downarrow 7.8Hz) and 8.28 (3, \downarrow 8.1Hz) together (1H, NHCO), 7.43-7.25 (5H, m, Ph), 7.14-7.08 (2H, m, Ar_H), 6.92-6.88 (2H, m, Ar_H), 5.06 (2H, s, CH₂Ph), 4.79-4.66 (m) and 4.45-4.42
35 (m) and 4.21 (d, \downarrow 9.7Hz) together (4H, CH α -tyr, CH α -thioprop and NCH₂S), 3.62 (s) and 3.61 (s) together (3H, CO₂Me), 3.28-2.71 (4H, m,

CHCH₂Ar + CHCH₂S) and 2.04 (s) and 1.82 (s) togher (3H, CH₃CO); m/z (ESI, 15V) 443 ($M^+ + 1$).

EXAMPLE 6

5 N-Acetyl-D-thiopropine-(O-benzyl)-L-tyrosine

Lithium hydroxide (47mg, 1.1mmol) was added to a solution of the compound of Example 5 (410mg, 0.93mmol) in a mixture of THF (10ml) and water 10ml). The mixture was stirred for 30min at room temprature then the THF was evaporated *in vacuo*. The aqueous residue was
 10 acidified (1M hydrochloric acid and extracted with DCM (2 x 50ml). The combined organic extracts were dried (Na₂SO₄) and evaporated *in vacuo* to give a gummy solid. This was dissolved in methanol, diluted with water and freeze-dried to give the title compound as a fluffy white solid (369mg, 93%). δ H (DMSO-d₆, 400K) 7.65 (1H, br d, CONH), 7.43-7.29 (5H, m,
 15 Ph), 7.12 (2H, d, \perp 8.5Hz, ArH), 6.91 (2H, d, \perp 8.6, ArH), 5.08 (2H, s, CH₂Ph), 4.81 (1H, dd, \perp 3.9-7.4Hz, CH α -thioprop), 4.76 (1H, d, \perp 9.2Hz, NCH_AH_BS), 4.50 (1H, dt, \perp 5.4, 8.3Hz, CH α tyr), 4.36 (1H, d, \perp 9.1Hz, NCH_AH_BS), 3.23 (1H, dd, \perp 7.1, 11.5Hz, CHCH_AH_BS), 3.07 (1H, dd, \perp 5.4,
 20 14.1 Hz, CH_AH_BAr), 2.99 (1H, dd, \perp 3.9, 11.5Hz, CHCH_AH_BS), 2.91 (1H, dd, \perp 8.4, 14.2Hz, CH_AH_BAr) and 1.97 (3H, s, CH₃CO) [COOH not observed at 400K. δ H (DMSO-d₆, 300K) 12.7 1Hv. br s. CO₂H]; m/z (ESI, 15V) 429 ($M^+ + 1$).

EXAMPLE 7

25 N-Acetyl-D-thiopropine-(O-2,6-dicholorobenzyl)-L-tyrosine methyl ester

A solution of Intermediate 2 (352mg, 1mmol) in DMF (5ml) was added to a suspension of sodium hydride (60% in mineral oil, 44mg, 1.1mmol) in DMF (3ml) at room temperature. After 5min a solution of 2,6 dichloro-benzyl
 30 bromide (288mg, 1.2mmol) in DMF (2ml) was added and the mixture stirred for 90min. The reaction was quenched with a few drops of water and the DMF was removed *in vacuo*. The residue was dissolved in ethyl acetate (100ml), washed with water (2 x 50ml) and brine (25ml), dried (Na₂SO₄) and evaporated *in vacuo*. The residue was purified by column chromatorgaphy (SiO₂; methanol/DCM 5:95) to give the title compound as
 35 a white foam (499mg, 97%); δ H (DMSO-d₆, 300K) (2 rotameric species

observed) 8.57 (d, J 8.2Hz) and 8.31 (d, J 8.1Hz), together (1H, NHCO), 7.57-7.43 (3H, m, OCH_2ArH), 7.18-7.12 (2H, t, J 8.1Hz, CHCH_2ArH), 6.98-6.94 (2H, m, CHCH_2ArH), 5.19 (2H, s, OCH_2Ar), 4.79-4.68 (m) and 4.50-4.44 (m) and 4.45 (d, J 8.8Hz), and 4.23 (d, J 9.7Hz) together (4H, 2 x $\text{CH}_\alpha + \text{NCH}_2\text{S}$), 3.64 (s) and 3.635 (s) together (3H, CO_2CH_3), 3.35-3.27 (m) and 3.17-2.49 (m) together (4H, $\text{CHCH}_2\text{Ar} + \text{CHCH}_2\text{S}$) and 2.05 (s) and 1.83 (s) together (3H, CH_3CON); m/z (ESI, 15V) 511 ($\text{M}^+ + 1$).

EXAMPLE 8

10 *N*-Acetyl-*D*-thiopropine-(*O*-2,6-dichlorobenzyl)-*L*-tyrosine

Lithium hydroxide (44mg, 1.05mmol) was added to a solution of the compound of Example 7 (490mg, 0.959mmol) in a mixture of THF (10ml) and water (10ml). The mixture was stirred at room temperature for 15min then the THF was evaporated *in vacuo*. The aqueous residue was
 15 acidified (1M hydrochloric acid) and extracted with DCM (2x50ml). The combined extracts were dried (Na_2SO_4) and evaporated *in vacuo* to give the title compound as a white solid (464mg, 97%) (Found: C, 53.02; H, 4.46; N, 5.50. $\text{C}_{22}\text{H}_{22}\text{N}_2\text{O}_5\text{SCl}_2$ requires C, 53.12; H, 4.46; N, 5.63%); δH (DMSO- d_6 , 400K) 7.63 (1H, brd, CONH), 7.51-7.38 (3H, m, OCH_2ArH),
 20 7.16 (2H, d, J 8.7Hz, CHCH_2ArH), 6.95 (2H, d, J 8.7Hz, CHCH_2ArH), 5.26 (2H, s, OCH_2Ar), 4.83 (1H, dd, 3.9, 7.4, $\text{CH}_\alpha\text{thioprop}$), 4.77 (1H, d, J 9.2Hz, $\text{NCH}_\text{A}\text{H}_\text{BS}$), 4.53 (1H, dt, J 5.4, 8.3Hz, $\text{CH}_\alpha\text{tyr}$), 4.37 (1H, d, J 9.2Hz, $\text{NCH}_\text{A}\text{H}_\text{BS}$), 3.24 (1H, dd, J 7.3, 11.5Hz, $\text{CHCH}_\text{A}\text{H}_\text{BS}$), 3.10 (1H, dd, J 5.4, 14.1Hz, $\text{CHCH}_\text{A}\text{H}_\text{BAr}$), 3.00 (1H, dd, J 3.9, 11.5 Hz, $\text{CHCH}_\text{A}\text{H}_\text{BS}$), 2.94
 25 (1H, dd J 8.4, 14.1 Hz, $\text{CHCH}_\text{A}\text{H}_\text{BAr}$) and 1.99 (3H, s, CH_3CO); m/z (ESI, 27V) 497 ($\text{M}^+ + 1$).

EXAMPLE 9

30 *N*-Acetyl-*D*-thiopropine-(*O*-2,6-dichlorobenzyl)-3-nitro-*L*-tyrosine methyl ester

A solution of Intermediate 4 (596mg, 1.5mmol) in DMF (5ml) was added to a suspension of sodium hydride (60% in mineral oil, 66mg, 1.65mmol) in DMF (10ml) at 0°. After 10min a solution of 2,6-dichlorobenzylbromide (432mg, 1.8mmol) in DMF (3ml) was added and the mixture stirred at 0°
 35 for 2h and at room temperature for 1h. The reaction was quenched with a few drops of water and the DMF removed *in vacuo*. The residue was

dissolved in ethyl acetate (100ml), washed with water (2x30ml) and brine (30ml), dried (Na_2SO_4) and evaporated *in vacuo*. The residue was purified by column chromatography (SiO_2 ; methanol/DCM 5:95) to give the title compound as a yellow oil (600mg, 72%); δH ($\text{DMSO}-d_6$, 300K) (2 rotameric species observed) 8.63 (d, J 8.0Hz) and 8.37 (d, J 8.3Hz) together (1H, CONH), 7.75-7.71 (1H, m, CHCH_2ArH), 7.58-7.45 (5H, m, ArH), 5.38 (2H, s, OCH_2Ar), 4.75-4.68 (m) and 4.56-4.48 (m) and 4.44 (d, J 8.6Hz) and 4.20 (d, J 9.7Hz) together (4H, $2\times\text{CH}_\alpha + \text{NCH}_2\text{S}$), 3.66 (3H, s, CO_2CH_3), 3.29-2.73 (4H, m, $\text{CHCH}_2\text{Ar} + \text{CHCH}_2\text{S}$) and 2.04 (s) and 1.85 (s) together (3H, CH_3CO); m/z (ES, 15V) 556 ($M^+ + 1$).

EXAMPLE 10

N-Acetyl-*D*-thiopropine-(*O*-2,6-dichlorobenzyl)-3-3 nitro-*L* tyrosine

Lithium hydroxide (49mg, 1.17mmol) was added to a solution of the compound of Example 9 (590mg, 1.06mmol) in a mixture of THF (10ml) and water (10ml). The mixture was stirred at room temperature for 30min then the THF was evaporated *in vacuo*. The aqueous residue was acidified (1M, hydrochloric acid) and extracted with ethyl acetate (2x75ml). The combined extracts were dried (Na_2SO_4) and evaporated *in vacuo*. The residue was freeze dried from a mixture of methanol and water to give the title compound as a fluffy yellow solid (510mg, 89%) (Found: C, 48.39; H, 3.91; N, 7.60. $\text{C}_{22}\text{H}_{21}\text{N}_3\text{O}_7\text{SCl}_2$ requires C, 48.72; H, 3.90; N, 7.75%); δH ($\text{DMSO}-d_6$, 400K) 7.85 (1H, br d, CONH), 7.67 (1H, d, J 2.2Hz, ArH), 7.54-7.41 (5H, m, ArH), 5.44 (2H, s, OCH_2Ar), 4.82 (1H, dd, J 4.0, 7.3Hz, $\text{CH}_\alpha\text{thioprop}$), 4.76 (1H, d, J 9.1Hz, $\text{NCH}_\text{A}\text{H}_\text{BS}$), 4.57 (1H, dt, J 5.2, 8.7Hz, $\text{CH}_\alpha\text{tyr}$), 4.37 (1H, d, J 9.2Hz, $\text{NCH}_\text{A}\text{H}_\text{BS}$), 3.28-3.16 (2H, m, $\text{CHCH}_\text{A}\text{H}_\text{BAr} + \text{CHCH}_\text{A}\text{CH}_\text{BS}$), 3.06-2.98 (2H, m, $\text{CHCH}_\text{A}\text{H}_\text{BAr} + \text{CHCH}_\text{A}\text{H}_\text{BS}$) and 1.99 (3H, s, CH_3CO); m/z (ES, 30V) 542 ($M^+ + 1$).

EXAMPLE 11

N-Acetyl-*O*-thiopropine- α -methyl-(*O*-2,6-dichlorobenzoyl)-*L*-tyrosine methyl ester

A solution of Intermediate 7 (347mg, 0.948mmol) in DMF (5ml) was added to a suspension of sodium hydride (60% in mineral oil, 40mg, 0.995mmol) in DMF (5ml) at room temperature. After 10min 2,6-dichlorobenzoyl chloride (150 μl , 1.04mmol) was added and the mixture stirred for 1h. The

reaction was quenched with a few drops of water and the DMF was evaporated *in vacuo*. The residue was dissolved in ethyl acetate (10ml), washed with water (2x30ml) and brine (30ml), dried (Na₂SO₄) and evaporated *in vacuo*. The residue was purified by column chromatography (SiO₂, methanol/DCM 5:95) to give the title compound as a yellow gum (423mg, 83%); δ H (DMSO-d₆, 400K) 7.63-7.46 (4H, m, CONH + COArH), 7.30 (2H, d, \downarrow 8.8Hz, ArH), 7.21 (2H, d, \downarrow 8.7Hz, ArH), 4.86 (1H, dd, \downarrow 3.89, 7.4Hz, CH α thioprop), 4.78 (1H, d, \downarrow 9.1Hz, NCH_AH_BS), 4.44 (1H, d, \downarrow 9.1Hz, NCH_AH_BS), 3.66 (3H, s, CO₂CH₃), 3.38-3.14 (4H, m, CHCH₂S + CH₂Ar), 2.05 (3H, s, CH₃CO) and 1.40 (3H, s, CCH₃); m/z (ES, 15V) 539 ($M^+ + 1$).

EXAMPLE 12

N-Acetyl-D-thiopropine- α -methyl-(O-2,6-dichlorobenzoyl)-L-tyrosine

Lithium hydroxide (34mg, 0.82mmol) was added to a solution of the Compound of Example 11 in a mixture of THF (7ml) and water (7ml). The mixture was stirred at room temperature for 7h and the THF was evaporated *in vacuo*. The aqueous residue was acidified (1M, hydrochloric acid) and extracted with DCM (2x75ml). The extracts were dried (Na₂SO₄) and evaporated *in vacuo*. Purification by column chromatography (SiO₂, methanol / acetic acid / DCM, 5:5:90) and freeze drying from a mixture of methanol and water gave the title compound as a fluffy white solid (275mg, 64%). (Found: C, 51.81; H, 4.18; N, 5.13. C₂₃H₂₂N₂O₆SCl₂, (H₂O)_{0.4} requires C, 51.87; H, 4.32; N, 5.26%); δ H (DMSO-d₆, 400K) 7.63-7.54 (3H, m, COArH), 7.43 (1H, br s, CONH), 7.31 (2H, d, \downarrow 8.7Hz, ArH), 7.19 (2H, d, \downarrow 8.7Hz, ArH), 4.86 (1H, dd, \downarrow 3.9, 7.4Hz, CH α thioprop), 4.79 (1H, d, \downarrow 9.2Hz, NCH_AH_BS), 4.42 (1H, d, \downarrow 9.2Hz, NCH_AH_BS), 3.38-3.26 (3H, m, CH_AH_BAr + CHCH_AH_BS), 3.17 (1H, dd, \downarrow 3.9, 11.5Hz, CHCH_AH_BS), 2.02 (3H, s, CH₃CO) and 1.45 (3H, s, CCH₃); m/z (ES, 15V) 525 ($M^+ + 1$).

EXAMPLE 13

N-Acetyl-D-thiopropine-(O-2,4,6-trichlorobenzyl)-L-tyrosine methyl ester

A solution of Intermediate 2 (528mg, 1.5mmol) in DMF (5ml) was added to a suspension of sodium hydride (60% in mineral oil, 66mg, 1.65mmol) in

DMF (5ml) at 0°. After 10 min a solution of Intermediate 8 (453mg, 1.65mmol) in DMF (5ml) was added and the mixture stirred at 0° for 2h. The reaction was quenched with a few drops of water and the DMF was removed *in vacuo*. The residue was dissolved in ethyl acetate (150ml), washed with water (2x50ml) and brine (25ml), dried (Na₂SO₄) and evaporated *in vacuo*. The residue was purified by column chromatography (SiO₂, ethyl acetate/hexane, 90/10) to give the title compound as a colourless viscous oil (619mg, 76%); δ H (DMSO-d₆, 300K) (2 rotameric species observed) 8.57 (d, J 8.2Hz) and 8.31 (d, J 8.2Hz) together (1H, CONH), 7.77 (2H, s, OCH₂ArH), 7.15 (2H, t, J 8.0Hz, CHCH₂ArH), 6.97-6.93 (2H, m, CHCH₂ArH), 5.15 (2H, s, OCH₂Ar), 4.79-4.67 (m) and 4.50-4.43 (m) and 4.46 (d, J 8.7Hz) and 4.22 (d, J 9.7Hz) together (4H, 2xCH α + NCH₂S), 3.63 (s) and 3.64 (s) together (3H, CO₂CH₃), 3.31-2.69 (4H, m, CHCH₂Ar + CHCH₂S) and 2.05 (s) and 1.83 (s) together (3H, CH₃CO); m/z (ES, 60V) 545 (M^+ + 1).

EXAMPLE 14

N-Acetyl-D-thioprolin-(O-2,4,6-trichlorobenzyl)-L-tyrosine

Lithium hydroxide (52mg, 1.23mmol) was added to a solution of the compound of Example 13 (610mg, 1.12mmol) in a mixture of THF (11ml) and water (11ml). The mixture was stirred at room temperature for 30min and the THF removed *in vacuo*. The aqueous residue was acidified (1M hydrochloric acid) and extracted with DCM (2x50ml). The combined extracts were dried (Na₂SO₄) and evaporated *in vacuo*. The residue was freeze dried from a mixture of methanol and water to give the title compound as a fluffy white solid (538mg, 90%). (Found: C, 49.17; H, 3.99; N, 5.18. C₂₂H₂₁N₂O₅SCl₃ · 0.25 (H₂O) requires C, 49.27; H, 4.04; N, 5.22%); δ H (DMSO-d₆, 400K) 7.65 (1H, br d, CONH), 7.61 (2H, s, OCH₂ArH), 7.16 (2H, d, J 8.7Hz, CHCH₂ArH), 6.95 (2H, d, J 8.7Hz, CHCH₂ArH), 5.23 (2H, s, OCH₂Ar), 4.82 (1H, dd, J 3.9, 7.4Hz, CH α thiopro), 4.77 (1H, d, J 9.2Hz, NCH_AH_BS), 4.52 (1H, dt, J 5.4, 8.3Hz, CH α tyr), 4.37 (1H, d, J 9.2Hz, NCH_AH_BS), 3.24 (1H, dd, J 7.4, 11.5Hz, CHCH_AH_BS), 3.10 (1H, dd, J 5.4, 14.2Hz, CHCH_AH_BAr), 3.00 (1H, dd, J 3.9, 11.5Hz, CHCH_AH_BS), 2.93 (1H, dd, J 8.4, 14.2Hz, CHCH_AH_BAr) and 1.98 (3H, s, CH₃CO); m/z (ES, 60V) 531 (M^+ + 1).

EXAMPLE 15**N-Acetyl-D-thiopropine-(O-2,6-difluorobenzyl)-L-tyrosine methyl ester**

Caesium carbonate (0.609g, 1.87mmol) was added in one portion to a solution of Intermediate 2 (0.60g, 1.70mmol) in DMF (15ml). α -Bromo-2,6-difluorotoluene (0.387g, 1.87mmol) was then added and the reaction stirred for 16h at room temperature. The reaction was partitioned between ethyl acetate (50ml) and water (30ml). The organic layer was separated and the aqueous layer was extracted with ethyl acetate (2x50ml). The combined organic layers were washed with brine (50ml), dried over MgSO_4 and the solvent removed under vacuum to give a white, waxy solid (1.0g). Trituration with diisopropyl ether gave a white solid which was isolated by filtration, washed with diisopropyl ether and dried to give the title compound (0.69g, 85%). δH (DMSO- d_6 , 390K) 7.84 (1H, br d, CONH), 7.52-7.42 (1H, m, OCH_2ArH), 7.15-7.06 (4H, m, $\text{OCH}_2\text{ArH} + \text{CHCH}_2\text{ArH}$), 6.93 (2H, d, J 8.4Hz, CHCH_2ArH), 5.13 (2H, s, OCH_2Ar), 4.82 (1H, dd, J 3.9, 7.2Hz, $\text{CH}_\alpha\text{thioprop}$), 4.76 (1H, d, J 9.2Hz, $\text{NCH}_\text{A}\text{H}_\text{BS}$), 4.56 (1H, dt, J 5.8, 8.2Hz, $\text{CH}_\alpha\text{tyr}$), 4.38 (1H, d, J 9.3Hz, $\text{NCH}_\text{A}\text{H}_\text{BS}$), 3.65 (3H, s, CO_3CH_3), 3.25 (1H, dd, J 7.4, 11.4Hz, $\text{CHCH}_\text{A}\text{H}_\text{BS}$), 3.10-2.91 (3H, m, $\text{CHCH}_2\text{Ar} + \text{CHCH}_\text{A}\text{H}_\text{BS}$) and 1.99 (3H, s, CH_3CO); m/z (ES) 479 ($\text{M}^+ + 1$).

EXAMPLE 16**N-Acetyl-D-thiopropine-(O-2,6-difluorobenzyl)-L-tyrosine**

A solution of the compound of Example 15 (0.69g, 1.44mmol) in dioxane/methanol (1:1, 8ml) and water (6ml) was treated with lithium hydroxide monohydrate (72.6mg, 1.73mmol) in one portion. A further 1ml of dioxane was added to give a clear solution. The reaction was stirred for 1.5h at room temperature and then acidified to pH 4.5 with a few drops of glacial acetic acid to give a whiter precipitate. The bulk of solvent was removed *in vacuo*, water (5ml) added and the solid isolated by filtration, washed well with water (20ml), hexane (20ml) and dried under vacuum to give the title compound as a white solid (0.56g, 96%). δH (DMSO- d_6 , 390K) 7.69 (1H, br d, CONH), 7.53-7.42 (1H, m, OCH_2ArH), 7.16-7.06 (4H, m, $\text{OCH}_2\text{ArH} + \text{CHCH}_2\text{ArH}$), 6.93 (2H, d, J 8.6Hz, CHCH_2ArH), 5.12 (2H, s, OCH_2Ar), 4.82 (1H, dd, J 3.9, 7.3Hz, $\text{CH}_\alpha\text{thioprop}$), 4.77 (1H, d, J 9.2Hz, $\text{NCH}_\text{A}\text{H}_\text{BS}$), 4.50 (1H, dt, J 5.4, 8.3Hz, $\text{CH}_\alpha\text{tyr}$), 4.37 (1H, d, J

9.0Hz, NCH_AH_BS), 3.24 (1H, dd, J 7.4, 11.5Hz, CHCH_AH_BS), 3.08 (1H, dd, J 5.4, 14.1Hz, CHCH_AH_BAr), 2.99 (1H, dd, J 3.9, 11.5Hz, CHCH_AH_BS), 2.92 (1H, dd, J 8.5, 14.2Hz, CHCH_AH_BAr) and 1.98 (3H, s, CH₃CO); m/z (ES) 4.65 ($M^+ + 1$).

5

The following assays can be used to demonstrate the potency and selectivity of the compounds according to the invention. In each of these assays an IC₅₀ value was determined for each test compound and represents the concentration of compound necessary to achieve 50% inhibition of cell adhesion where 100% = adhesion assessed in the absence of the test compound and 0% = absorbance in wells that did not receive cells.

$\alpha_4\beta_1$ Integrin-dependent Jurkat cell adhesion to VCAM-Ig

96 well NUNC plates were coated with F(ab)₂ fragment goat anti-human IgG Fc γ -specific antibody [Jackson Immuno Research 109-006-098: 100 μ l at 2 μ g/ml in 0.1M NaHCO₃, pH 8.4], overnight at 4°. The plates were washed (3x) in phosphate-buffered saline (PBS) and then blocked for 1h in PBS/1% BSA at room temperature on a rocking platform. After washing (3x in PBS) 9 ng/ml of purified 2d VCAM-Ig diluted in PBS/1% BSA was added and the plates left for 60 minutes at room temperature on a rocking platform. The plates were washed (3x in PBS) and the assay then performed at 37° for 30 min in a total volume of 200 μ l containing 2.5 x 10⁵ Jurkat cells in the presence or absence of titrated test compounds.

25

Each plate was washed (2x) with medium and the adherent cells were fixed with 100 μ l methanol for 10 minutes followed by another wash. 100 μ l 0.25% Rose Bengal (Sigma R4507) in PBS was added for 5 minutes at room temperature and the plates washed (3x) in PBS. 100 μ l 50% (v/v) ethanol in PBS was added and the plates left for 60min after which the absorbance (570nm) was measured.

30

$\alpha_4\beta_7$ Integrin-dependent JY cell adhesion to MAdCAM-Ig

This assay was performed in the same manner as the $\alpha_4\beta_1$ assay except that MAdCAM-Ig (150ng/ml) was used in place of 2d VCAM-Ig and a sub-line of the β -lympho blastoid cell-line JY was used in place of Jurkat cells.

35

The IC₅₀ value for each test compound was determined as described in the $\alpha_4\beta_1$ integrin assay.

$\alpha_5\beta_1$ Integrin-dependent K562 cell adhesion to fibronectin

- 5 96 well tissue culture plates were coated with human plasma fibronectin (Sigma F0895) at 5 μ g/ml in phosphate-buffered saline (PBS) for 2 hr at 37°C. The plates were washed (3x in PBS) and then blocked for 1h in 100 μ l PBS/1% BSA at room temperature on a rocking platform. The blocked plates were washed (3x in PBS) and the assay then performed at 10 37°C in a total volume of 200 μ l containing 2.5x 10⁵ K562 cells, phorbol-12-myristate-13-acetate at 10ng/ml, and in the presence or absence of titrated test compounds. Incubation time was 30 minutes. Each plate was fixed and stained as described in the $\alpha_4\beta_1$ assay above.

15 **$\alpha_m\beta_2$ -dependent human polymorphonuclear neutrophils adhesion to plastic**

- 96 well tissue culture plates were coated with RPMI 1640/10% FCS for 2h at 37°C. 2 x 10⁵ freshly isolated human venous polymorphonuclear neutrophils (PMN) were added to the wells in a total volume of 200 μ l in the presence of 10ng/ml phorbol-12-myristate-13-acetate, and in the presence or absence of test compounds, and incubated for 20min at 37°C followed by 30min at room temperature. The plates were washed in medium and 100 μ l 0.1% (w/v) HMB (hexadecyl trimethyl ammonium bromide, Sigma H5882) in 0.05M potassium phosphate buffer, pH 6.0 added to each well. 20 The plates were then left on a rocker at room temperature for 60 min. Endogenous peroxidase activity was then assessed using tetramethyl benzidine (TMB) as follows: PMN lysate samples mixed with 0.22% H₂O₂ (Sigma) and 50 μ g/ml TMB (Boehringer Mannheim) in 0.1M sodium acetate/citrate buffer, pH 6.0 and absorbance measured at 630nm. 25

30

α_{IIb}/β_3 -dependent human platelet aggregation

- Human platelet aggregation was assessed using impedance aggregation on the Chronolog Whole Blood Lumiaggregometer. Human platelet-rich plasma (PRP) was obtained by spinning fresh human venous blood 35 anticoagulated with 0.38% (v/v) tri-sodium citrate at 220xg for 10 min and

diluted to a cell density of $6 \times 10^8/\text{ml}$ in autologous plasma. Cuvettes contained equal volumes of PRP and filtered Tyrode's buffer (g/liter: NaCl 8.0; $\text{MgCl}_2 \cdot \text{H}_2\text{O}$ 0.427; CaCl_2 0.2; KCl 0.2; D-glucose 1.0; NaHCO_3 1.0; $\text{NaHPO}_4 \cdot 2\text{H}_2\text{O}$ 0.065). Aggregation was monitored following addition of
5 2.5 μM ADP (Sigma) in the presence or absence of inhibitors.

In the above assays the compounds of the invention generally have IC_{50} values in the $\alpha_4\beta_1$ and $\alpha_4\beta_7$ assays of 1 μM and below. The compounds of the Examples typically had IC_{50} values of 500nM and below in these
10 assays. In the other assays featuring α integrins of other subgroups the same compounds had IC_{50} values of 50 μM and above thus demonstrating the potency and selectivity of their action against α_4 integrins.